

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|--|--|---|---|
| (51) International Patent Classification ⁶ : C07K 14/085, A61K 39/125, G01N 33/569 | | A1 | (11) International Publication Number: WO 98/11133 |
| | | | (43) International Publication Date: 19 March 1998 (19.03.98) |
| (21) International Application Number: PCT/SE97/01515 | | (81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). | |
| (22) International Filing Date: 9 September 1997 (09.09.97) | | | |
| (30) Priority Data: ✓ 9603305-5 11 September 1996 (11.09.96) SE | | | |
| (71)(72) Applicant and Inventor: NIKLASSON, Bo [SE/SE]; Sibyllegatan 15, S-114 42 Stockholm (SE). | | Published With international search report. | |
| (74) Agents: NILSSON, Brita et al.; Oscar Grahn Patentbyrå AB, P.O. Box 19540, S-104 32 Stockholm (SE). | | | |

(54) Title: NEW PICORNAVIRUSES, VACCINES AND DIAGNOSTIC KITS

SEQ ID NO: 1 (Ljungan 87-012)

```
AGTCTAGTCT TATCTTGAT GTGTCCTGCA CTGAACCTGT TTCTGTCTCT 50
GGAGTGCTCT AACTTCAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC 100
ACAGGAATCT GCACAGGTGG CTTTCACCTC TGGACAGTGC ATTCCACACC 150 (1)
CGCTCCACGG TAGAAGATGA TGTGTGTCTT TGCTTGTAAG AAGCTTGTGA 200
AAATCGTGTG TAGGCGTAGC GGCTACTTGA GTGCCAGCGG ATTACCCCTA 250
GTGGTAACAC TAGC
```

(57) Abstract

A new group of picornaviruses is disclosed. The picornaviruses of the invention comprise in the non-coding region of their viral genome a nucleotide sequence which corresponds to cDNA sequence (I) or homologous sequences having at least 75 % homology to the SEQ ID NO:1, and they cause mammalian disease. Further aspects of the invention comprise a protein corresponding to a protein of the picornaviruses, antiserum or antibody directed against a protein of the picornaviruses, antigen comprising a protein of the picornaviruses, diagnostic kits, vaccines, use of the picornaviruses in medicaments, particularly for the treatment or prevention of Myocarditis, Cardiomyopathia, Guillain Barré Syndrome, and Diabetes Mellitus, Multiple Sclerosis, Chronic Fatigue Syndrome, Myasthenia Gravis, Amyotrophic Lateral Sclerosis, Dermatomyositis, Polymyositis, Spontaneous Abortion, and Sudden Infant Death Syndrome, and methods of treatment of diseases caused by the picornaviruses.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | | | PL | Poland | | |
| CN | China | KR | Republic of Korea | PT | Portugal | | |
| CU | Cuba | KZ | Kazakstan | RO | Romania | | |
| CZ | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | |
| DE | Germany | LI | Liechtenstein | SD | Sudan | | |
| DK | Denmark | LK | Sri Lanka | SE | Sweden | | |
| EE | Estonia | LR | Liberia | SG | Singapore | | |

NEW PICORNAVIRUSES, VACCINES AND DIAGNOSTIC KITS.5 **FIELD OF INVENTION**

The present invention relates to new picornaviruses, proteins expressed by the viruses, antisera and antibodies directed against said viruses, antigens comprising structural proteins of said viruses, diagnostic kits, vaccines, use of said viruses, antisera or antibodies and antigens in medicaments, and methods
10 of treating or preventing diseases caused by said viruses, such as Myocarditis, Cardiomyopathia, Guillain Barré Syndrome, and Diabetes Mellitus, Multiple Sclerosis, Chronic Fatigue Syndrome, Myasthenia Gravis, Amyotrophic Lateral Sclerosis, Dermatomyositis, Polymyositis, Spontaneous Abortion, and Sudden Infant Death Syndrome.

15

BACKGROUND OF THE INVENTION

Recently, a sudden death syndrome among Swedish orienteers has been observed. Of approximately 200 elite orienteers six died in myocarditis during 1989-1992 (1). Orienteering, aiming to find the fastest/shortest way between
20 several checkpoints and often in forested areas, is exceptional with respect to environmental exposure. Thus it has been speculated, that the sudden deaths syndrome among orienteers is caused by a vector borne (rodent or arthropod) infectious agent.

25 It has now been shown in an epidemiological study that the incidence of deaths in myocarditis in northern Sweden tracked the 3-4 year population fluctuations (cycles) of bank voles (*Clethrionomys glareolus*) with one year time lag. Previously, it has been shown that cardioviruses, with rodents as their natural reservoir, can cause Guillain Barré Syndrome (GBS) in man, Diabetes Mellitus
30 (DM) in mice and myocarditis in several species including non-human primates.

In addition to death in myocarditis it is also shown in the epidemiological study that the number of patients diagnosed with Guillain Barré Syndrome (GBS), and Diabetes Mellitus (DM) in north m Swed n tracked the 3-4 year population fluctuations of bank voles with different time delays.

5

Sven Gard and co-workers studied antibody prevalence to encephalomyelitis virus (EMCV) in Swedish normal population in the early 1950th (2). These studies found a surprisingly high antibody prevalence rate by hemagglutination inhibition test but no sera could be confirmed by neutralization test. These results were found puzzling at the time but could be explained by the presence of one or several related picornaviruses circulating in Sweden.

10

The fact that enterovirus have a large number of members and cardiovirus only two possibly three could reflect the true diversity of the two genus or only be the result of the amount of effort made to isolate new viruses from rodents as compared to isolating new enteroviruses from humans.

15

The Picornavirus family is presently divided into five genera (aphto-, entero-, hepato-, rhino-, and cardioviruses) (3). This taxonomy was initially based on morphological, physiological and serological properties as well as on the pathogenicity of the viruses. More recently, however, viruses have been characterized based on their genome sequence since it has been established that sequence data to a large extent coincide with the characterisation properties used previously (4,5).

20

25

The prototype virus in the cardiovirus genus is Theiler's murine encephalomyelitis virus (TMEV). Another member in this genus is encephalomyocarditis virus (EMCV). Vilyuisk virus, isolated from patients in Russia with degenerative neurological disease, is serologically related to TMEV but presently under consideration for being included as a third distinct member of the cardiovirus genus (6).

30

In nature, cardioviruses have a geographically widespread distribution and a large number of susceptible hosts with rodents as their natural reservoir. In addition to rodents, EMCV has been isolated from domestic pigs, elephants, lions, non human primates and man (7,8,9). Infection with TMEV and EMCV have provided excellent animal models for inducing myocarditis, DM and different neurological disorders such as demyelinating diseases resembling multiple sclerosis in mice (10-16). Other neurological or muscular disorders in which an infection is suspected to be the triggering factor and in which there is also an autoimmune component are Cardiomyopathia, Multiple Sclerosis (MS), Chronic Fatigue Syndrome (CFS), Myasthenia Gravis (MG), and Amyotrophic Lateral Sclerosis (ALS). It has never been established, however, that cardiovirus is a significant human pathogen, as disease in man most often has been described in case reports or as infection measured in sero-epidemiological surveys (7-17).

Thus, there may be other not yet identified picornaviruses circulating in the wild rodent population and occasionally infecting humans resulting in Myocarditis, Cardiomyopathia, Guillain Barré Syndrome, and Diabetes Mellitus, Multiple Sclerosis, Chronic Fatigue Syndrome, Myasthenia Gravis, Amyotrophic Lateral Sclerosis, Dermatomyositis, Polymyositis, Spontaneous Abortion, and Sudden Infant Death Syndrome, in genetically susceptible individuals.

The epidemiological link between important human diseases and small rodent abundance and what is previously known about picornavirus/cardiovirus motivated attempts to isolate novel picornaviruses from small rodents.

DESCRIPTION OF EXPERIMENTAL WORK AS BASIS FOR THE INVENTION

Trapping of animals

Small rodents were trapped at several locations in northern Sweden and transported live to the Swedish Institute for Infectious Disease Control in Stockholm, Sweden. Species, date and location of trapped animals were

recorded. Animals were bled using the anaesthesia and killed. Organs were immediately removed and stored at -70°C until tested for presence of virus. A total of 53 *Clethrionomys glareolus* and 28 *Microtus agrestis* were tested for virus isolation.

5

Virus isolation

The isolation technique used in the present study was different from what is most often used. The cells used for isolation were kept for a minimum of two weeks and virus growths were detected by both CPE (cytopathogenic effect) and by staining the cells by a large number of human sera using IFT (immunofluorescence test). None of the new viruses presented herein would have been isolated using routine procedure for detecting

10

cardioviruses/picornaviruses. They grow to lower titer and CPE develops slowly. Saliva mixed with lung homogenate and faeces were analyzed separately from each animal. The material was inoculated into T25 flask of confluent BHK-21 cells. Cells were blind passed twice a week during two weeks. At the end of this period or earlier if signs of CPE occurred, cells were removed from the T25 flask by a rubber policeman, placed onto 10-well spot slides, air dried and acetone fixed. The cells were then stained with panels of human sera including 5 multiple sclerosis patients, 5 patients recently diagnosed with DM and 5 athletes dying in myocarditis and bled at autopsy. All T25 flasks (saliva-lung and faeces separately) were tested individually by IFT using the complete panel of human sera at a 1:10 dilution.

20

25

Cells showing positive reaction by IFT using the human serum panels were selected for further analysis. This included inoculation intracerebrally into 1 day old suckling mice, serological characterisation and sequence analysis.

Antisera and serological procedures

30

Antisera to the virus isolates were raised in mice (NMRI), and Guinea Pigs (Dunkin Hartley). The animals were injected with a cell culture supernatant from

(BHK-21 cells) intraperitoneally and serum collected 4-6 weeks later.

Preimmunization sera were tested individually while postimmunization sera were pooled from all infected animals.

5 An indirect immunofluorescence test (IFT), as described previously (18,19) was used to test antibody titres in immunized animals. Briefly, spot slides were prepared by incubating virus on Green Monkey Kidney (GMK) cells for 6-10 days. At sign of discrete CPE cells were removed from the flask by a rubber policeman and put onto the microscope slides, air dried, and fixed in cold (4°C) acetone and stored at -70°C. The titer was determined by incubating serum
10 diluted in PBS in the slides at 37°C for 1 hour in a moist chamber, followed by a FITC conjugate (F(ab')₂ fragment of goat anti human IgG γ -chain specific, Sigma Immuno Chemicals (F-1641) or Rabbit anti mouse immunoglobulins Daco (F0313)) incubated as above.

15 Antibody titers to the viruses were determined by a modification of the Plaque reduction neutralization test (PRNT) as described by Earley et al (20). In the test, sera were serially diluted four-fold and mixed with an equal volume containing 80-100 plaque-forming units (pfu) of virus per 50 μ l. The mixtures
20 were then incubated at 37°C for 60 minutes, and 50 μ l subsequently inoculated into each of 2 wells of a tissue culture plate containing confluent Vero cell monolayer. After adsorption for 60 minutes at 37°C the wells were overlaid with 0.5 ml of a 42°C mixture of 1 part 1% agarose and 1 part 2X basal Eagle's medium with Earle's salts, 17 mM Hepes buffer, 8% heated fetal calf serum, 100
25 U/ml penicillin, 100 μ g streptomycin. The tissue culture plates were incubated at 37°C in a humidified 5% CO₂ atmosphere for 3-7 days. A second overlay (0.5 ml) containing neutral red stain (1:9000) was then applied and plaques were enumerated the following day. The plaque numbers were linearly extrapolated to 2-fold dilutions. An 50% reduction of plaques was used as the criterion for
30 virus neutralization titers.

Electron microscopy

Cell culture media or brain tissue homogenates were examined by negative contrast electron microscopy (EM). A 10 µl droplet was incubated on Formvar/carbon-coated grids for one minute or alternatively, 0.5 ml samples were centrifuged for 30 minutes at 20,000 x g to remove cell debris and finally the supernatants were pelleted directly onto grids in a Beckman Airfuge for 10 minutes at 160,000 x g. Grids were stained with 2% phosphotungstate acid (pH 6.0) and examined in a Philips CM 100 electron microscope at a magnification of at least 46,000.

Sequence data

The isolates 87-012, 174F and 145SL were grown on the human lung carcinoma line A549 in 1600 cm² roller bottles. Full CPE was observed after 5-10 days. Supernate was filtered through 0.45 µm cellulose acetate filters (Costar) and the virus was pelleted at 20,000 g for 20 h at 4°C. RNA was isolated from the virus containing pellets using acid guanidinium thiocyanate as described (Chomczynski and Sacchi). Synthesis of cDNA was performed under standard conditions using 1 µg of RNA, AMV reverse transcriptase (Boehringer-Mannheim) and random 14 mer oligonucleotides as primers in a 20 µL reaction. Fragments of the viral 5'UTR were amplified using cardiovirus specific consensus primers: (sense) 5'-GGCCGAAGCCGCTTGAATA-3' (SEM) and (antisense) 5'-GTGGCTTTTGGCCGCAGAG-3' (ATVEM), both primers modified after the EMCV2 and EMCV1 primers previously reported (Jongen et al. 1993. Ann. Reum. Dis. 52:575-578. Cardiovirus sequences were from Dr A. Palmenberg (personal communication). Amplification conditions were 30 cycles at: 94°C, 30 sec., 50°C, 30 sec, 72°C, 2 min. The amplified fragments were cloned into the pCRII T-vector (In-Vitrogen). The cloned viral sequences were sequenced using A Taq polymerase FS cycle sequencing kit and data was collected on a ABI Prism 310 sequencing machine using M13 -21 and M13 reverse primers (Perkin-Elmer). A 1.8 kb fragment extending from the 5'-UTR into the viral polyprotein sequences was obtained by PCR (polymerase chain reaction) amplification of cDNA from the 145SL isolate. The primers were:

(sense) 5'-ACAGTGCATTCCACAC-3' (SLJU1) or 5'-CCGCTCCACAATAGA-3' (SLJU2) and (antisense) 5'-GATCTCAGAC-3' (primer 118). The SLJU1 and SLJU2 primers are located immediately adjacent to one another and were chosen as consensus primers for the Ljungan isolates of the invention with as little homology as possible to the EMCV and TMEV groups of viruses. The amplification conditions were 30 cycles at: 94°C, 30 sec., 42°C, 1 min, 72°C 2 min. The antisense primer 118 yielded similarly sized PCR products with either the SLJU1 or SLJU2 as sense primers, but none of the primers yielded PCR fragments when used alone. The sequence of the primer 118 was previously published (Bauer, D., et al. 1993. Nucl. Acids Res. 21:4272-4280). The obtained 1.8 kb PCR fragment was cloned and sequenced as described above.

RESULTS OF EXPERIMENTAL WORK

Three virus isolates were selected based on reaction with the human serum panels and showing a size and structure compatible with a picornavirus on EM. The first isolate was named Ljungan 87-012. Ljungan is a river in Medelpad county, Sweden where the animals were trapped.

The second and third isolate were designated Ljungan 174F and Ljungan 145SL, respectively.

All three isolates came from *C. glareolus*.

All three isolates killed suckling mice in 3-5 days.

The titer in mouse brain was 10^9 (approximately) while the cell culture titer was only 10^5 (approximately).

Electron microscopy

Virus particles, 27 nm in diameter, were spherical with the surface almost featureless and they appeared single or in small aggregates. In rare cases the stain penetrated the particles which made them look like empty shells.

Serological results

It was found after testing a number of different cell lines the Green Monkey Kidney cells were most suitable for making IFT drop slides for serology. The cross IFT data using mouse sera are seen in Table 1.

TABLE 1

Cross-IFT using virus infected GMK cells. Immune mice were titrated using 4 fold dilutions starting at a 1:10 dilution.

| | VIRUS | | |
|----------|--------|------|-------|
| | 87-012 | 174F | 145SL |
| Antisera | | | |
| 87-012 | 2560 | 160 | <10 |
| 174F | 160 | 160 | <10 |
| 145SL | 40 | 40 | 640 |

PRNT (plaque reduction neutralization test) data, preliminary results. Rabbit sera against TEMV and EMCV with a titer of 1:160 homologous had a titer less than 10 to the three isolates. Several attempts to make antisera with neutralizing titer in bank voles, mice, rabbits and guinea pigs have failed. All animals made high titer antibodies by IFT but no by PRNT. Bank voles failed to make IFT antibodies.

Sequence data

Sequences from 5'UTR and polyprotein gene of Ljungan virus isolates.

Cardiovirus consensus primers yielded a product of 303 bp for the three isolates 87-012, 174F and 145SL, compared to 284 bp for EMC virus. The fragment amplified was located immediately after the end of the poly C tract in EMC virus. PCR products specific for the Ljungan isolates were only obtained when the reannealing temperature was 50°C, and not at 58°C, which was optimal for

obtaining products from EMC virus cDNA. The subsequent sequence analysis revealed that the ATVEM primer was mismatched at 4 internal positions, explaining this difference in reannealing temperature. An alignment of the 5'UTR sequences for the three Ljungan isolates, EMCV and Vilyuisk virus (Table 2) shows a greater similarity between EMCV and Vilyuisk virus than between either of the two and the Ljungan isolates. It also demonstrates that each Ljungan isolate is distinct from the other by a number of nucleotide changes. The 174F and 145SL are similar to the isolate 87-012. The sequence homology between 174F and 87-012 was at most 95% (three undetermined bases in the sequence) while the homology between 87-012 and 145SL was 91%.

The strategy chosen for obtaining additional PCR fragments from the Ljungan virus isolates was a modification of a technique for detecting differentially expressed mRNAs (Bauer, D., et al. 1993. Nucl. Acids Res. 21:4272-4280). As a test for this strategy, cDNA from the Ljungan 145SL isolate was amplified using the conditions above, using either the SLJU1 or the SLJU2 primer as a sense primer and one of twenty 10-mer oligonucleotides of randomly chosen sequence as "antisense" primer.

If the PCR products obtained with the SLJU1 or SLJU2 primers and a specific 10-mer were similarly sized, and none of the primers yielded a product of this size when used alone in the PCR reaction, the fragment obtained was isolated and cloned. Only one combination of primers satisfied this criterion, namely the SLJU1 or SLJU2 primers in combination with the 118 10-mer oligonucleotide, which yielded a 1.8-1.9 kb PCR product. Of this fragment, 819 bp were sequenced from the 3' end. This sequence contained an open reading frame (ORF) of 663 bp in the sense of the viral polyprotein. This ORF was used to search in the Swiss protein data bank using the BLITZ search service from EMBL with the default search parameters. The top 10 scores were picornavirus polyprotein sequences, including 8 cardiovirus sequences. Homology was found over 188 a.a. The relatedness of this segment of the viral polyprotein to previously sequenced cardioviruses is shown in Table 3. A comparative

alignment of all cardioviruses was made available to us by Dr. A. Palmenberg. In Table 3, the sequence of TMEB An was arbitrarily taken as the index strain. For the 12 remaining cardioviruses in the alignment, only differences in amino acid sequence are shown. The alignment of the Ljungan 145SL sequence is similarly represented at the top. Since the BLITZ search algorithm takes into account identical as well as similar amino acids, the latter have been indicated by small type, while differences to TMEBeAn is in capitals as for the other strains in the alignment.

In conclusion, the above presented data for the Ljungan isolates are characteristic for the 3 viruses but yet incomplete. However, the comparison of cloned sequences from both a highly conserved part of the 5'-untranslated region of cardioviruses and coding sequences for the viral capsid proteins of one isolate (Ljungan 145SL) clearly show that the Ljungan viruses are related to the cardioviruses, but are more distant relatives than any previously identified cardiovirus. While the amino acid homology (identical amino acids) of the viruses within the Theiler group is 96-97%, the homology to Vilyuisk virus is about 83%, and the EMC viruses are 67-74% homologous to TMEBeAn, the Ljungan 145SL has only about 32% identical amino acids to TMEBeAn. Even if homology is taken as identical and similar amino acids, this measure of relationship would still amount to only 50% between Ljungan 145SL and TMEBeAn (the corresponding figure would be 79-83% between EMC and TMEBeAn).

ALIGNMENT OF SEQUENCES

Table 2 shows an alignment of three Ljungan virus isolates (1. 87-012, 2. 174F, 3. 145SL)[SEQ ID NO: 1,2 and 3, respectively] with published cardiovirus sequences (4. TMEBeAn, 5. Vilyuisk, 6. EMCV). The aligned sequence starts 29 nt 3' of the end of the poly-C tract in EMCV, and the sequence corresponds to nt 557 - 808 (approximately) in the different viral genomes. Inserted spaces in the sequences are indicated by a period (.).

TABLE 2

| | |
|----|--|
| 5 | 1. AGTCTAGTCTTATCTTGTATGTGTCCTGCACT..GA..ACTTGTTTCTGT 2. AGTCTAGTTTCATTCTGTGTGTGTTTGGCACT..GA..AATTATTTCTGT 3. AGTTTGGTTCTCTCTTGAGTGTGTTTTGTGTT..AG..CATAATTTCTGT 4. TGACAGG.GTTATTTTCACC.TCTTCTT..TTCTACTCCACAG.TG.T.T 5. TGACAGG.GTTATTTTCACC.TCTTCTCTCTTCTACTTCATAG.TG.T.T 6. AGGCCGGTGTGCGTTTGTCTATATGTTATTTTCCAC..CATA.TTGCCGT |
| 10 | 1. CTCTGGAGTGCTCTACACTTCAGTAGGGGCTGT.A.CCCGGGCGGTCCCA 2. CTCTGGGGTGCTTTACACTTCAGTAGGGGCTGT.A.CCCGGGCGGTCCCA 3. CTCTAGAGTGCTTTACACTCTAGTAGGGGCTGT.A.CCCGGGCGGTCCCA 4. CT.A.....TACTGTG..GAAGGGTATGTGT....TGCCCCTTCCT 5. CT.A.....TACTATG.AA.AGGGTATGTGT..C..GCCCTTCCT 6. CT.T.....TTGGCAATGT.G.AGGGCCCG.GAAACCTGGCCCTGTCT |
| 15 | 1. CTCTTCACAGGAATCTGCACAGGTGGCTTTTAC.CTCTGGACAGTGCATT 2. CTCTTCACAGGAATNTGCACAGGTGGCTTTTAC.CTCTGGACAGTGCATT 3. CTCTTCACAGGAATCTGCACAGGTGGCTTTTAC.CTCTGGACAGTGCATT 4. .TCTTGGAGAACGT..GCGCGGCGGTCTTTCCGTCTCTCGACAA.GCGC. 5. .TCTTGGAGAACGT..GCGTGGCGGTCTTTCCGTCTCTCGAAAAACG..T 6. .TCTTGACGAGCAT.T.CCTAGGGGTCTTTCCC.CTCTCGCCAAAGGAAT |
| 20 | 1. CCACACCCG.C.TCCACGGTAGAAGATGATGTGTGTCTTTGCT..TGTGA 2. CCACACCCG.C.TCCACAGTAGAAGATGATGTGTGTCTTTGCT..TGTGA 3. CCATACCCG.C.TCCACAATAGAAGATGATGTATATCTTTGTT..TGTGA 4. GCGT..GCAACATACAGAGT.AACG.CGAAGAA.AGCA..GTTT.CC.GG 5. GCGT..GCGACATGCAGAGT.AACG.CAAAGAA.AGCA..GTTT.T.TGG 6. GCA.A.G.GTC.TGTTGAAT.GTCG.TGAAGGA.AGCA..GTTTCTCTGG |
| 25 | 1. AAA.GCTT...GTGAAAATC.....GTGTGTAGGCGTAGCGGCTACT 2. AAA.GCTT...GTGAAAATC.....GTGTGTAGGCGTAGCGGNTACT 3. AAT.GCT.CA..TGAA.A.C.....GTGTGTGTAGGCGTAGCGGCTACT 4. TCTAGCT.CTAGTGCCCA.CAAGAAAACAGCTGTAG.CG.ACCA.C.ACA 5. TCTAGCT.CTGGTGCCCA.CAAGAAAACAGCTGTAG.CG.ACCA.C.ACA 6. AA..GCTTCT..TGAAGA.CAA.ACAACGTCTGTAG.CG.ACC..CT..T |
| 30 | 1. TGAGTGCCAGCGGATTACCCCTAGTGGTAACACTAGC 2. TGAGTGCCAGCGGACNACCCCTAGTGGTAACACTAGC 3. TGAATGCCAGCGGAACCCCCCTAGTGGTAACACTAGC 4. ..AAGGC.AGCGGAACCCCCCTCCTGGTAACAGGAGC 5. ..AAGGC.AGCGGAACCCCCCTCCTGGTAACAGGAGC 6. TGCAGGC.AGCGGAACCCCCCACCTGGCGACAGGTGC |
| 45 | In this region of the viral genome, Ljungan 174F has 94% homology to Ljungan 87-012 (here taken as the indicator strain for comparisons), and Ljungan 145SL |

has 91% homologous residues to Ljungan 87-012. The TMEBeAn strain has 69%, Vilyuisk has 68% and EMCV has 68% homologous residues to Ljungan 87-012. Using the same criteria for calculating the homology, EMCV has 85% homology to TMEBeAn.

5

Table 3 shows alignment of cDNA sequences from the polyprotein coding sequences of the Ljungan 145SL isolate [SEQ ID NO. 4] to the amino acid sequences of sequenced cardioviruses in the comparative alignment compiled by Dr. A. Palmenberg (personal comm.) The TMEBeAn strain was arbitrarily taken as the indicator strain, while the amino acids of the remaining strains are shown only if they differ from the indicator strain. For the Ljungan 145SL isolate, similar, but non-identical amino acids are indicated in small type. The amino acid homology between Ljungan 145SL and other cardioviruses was established screening the entire Swiss Protein Data Bank using the BLITZ search algorithm with standard search parameters.

10

15

TABLE 3

| | | | |
|----------|---|-----|-----|
| Ljungan | 464 | 480 | 525 |
| 145SL | K--m-iArm-sVyK-ERTEPGGTNG--QWshthSPInW-.fDGGiHLED-P-...-LFsSCy- | | |
| TMEBeAn | SDLLELCKLPT.FLGNPNTNNKRYPYFSATNSVPATSMVDYQVALSCSCMANSHLAAVARNFN | | |
| TMEGd7 | -----S-D-----L----- | | |
| TMEGd7 | -----S-D-----L----- | | |
| TMEDa | -----S-----T-L-----C----- | | |
| Vilyuisk | T-----L.S-DT-V-F-T-----TE-L-E--T-----S-----S----- | | |
| EMCBd | K-F--IAQI--.-I--KIP-AVP-IEA-N-.A-KTQPLAT--T-----L--TF--LS--A | | |
| EMCBc | K-F--IAQI--.-I--KIP-AVP-IEA-N-.A-KTQPLAT--T-----L--TF--LS--A | | |
| EMCDd | K-F--IAQI--.-I--KIP-AVP-IEA-N-.A-KTQPLAT--T-----L--TF--LS--A | | |
| EMCDc | K-F--IAQI--.-I--KIP-AVP-IEA-N-.A-KTQPLAT--T-----L--TF--LS--A | | |
| EMCDv1 | K-F--IAQI--.-I--KIP-AVP-IEA-N-.A-KTQPLAT--T-----L--TF--LS--A | | |
| EMCR | K-F--IAQI--.-I--KIP-AVP-IEA-N-.A-KTQPLAT--T-----L--TF--LS--A | | |
| MengoM | K-F--IAQI--.-I--KMP-AVP-IEA-N-.A-KTQPLAV--T-----L--TF--LS--A | | |
| Mengo37a | K-F--IAQI--.-I--KVP-AVP-IEA-N-.A-KTQPLAV--T-----L--TF--LS--A | | |

| | | | |
|----------|---|-----|-----|
| Ljungan | 526 | 540 | 588 |
| 145SL | Yw---TVLKLTVYAsTFN--rLRm-fF-I.MMqG-Q-.kKHkCLfMvC-i---nt-EM-I-y. | | |
| TMEBeAn | QYRGSNLNLFVFTGAAMVKGKFLIAYTPPGAGKPTTRDQAMQSTYAIWDLGLNSSFNFTAPFI | | |
| TMEGd7 | -----R-----A----- | | |
| TMEGd7b | -----A----- | | |
| TMEDa | -----A-----V----- | | |
| Vilyuisk | -----S--T-----X-----V----- | | |
| EMCBd | -----VYT-----T-M-----S-----A-----YS--V-- | | |
| EMCBc | -----VYT-----T-M-----S-----A-----YS--V-- | | |
| EMCDd | -----VYT-----T-M-----S-----A-----YS--V-- | | |
| EMCDc | -----VYT-----T-M-----S-----A-----YS--V-- | | |
| EMCDv1 | -----VYT-----T-M-----S-----A-----YS--V-- | | |
| EMCR | -----VYT-----T-M-----S-----A-----YS--V-- | | |
| MengoM | -----VYT-----T-M-----S-----A-----YS--V-- | | |
| Mengo37a | -----VYT-----T-M-----S-----A-----YS--V-- | | |

| | | | |
|----------|--|-----|-----|
| Ljungan | 589 | 600 | 651 |
| 145SL | ...w...GnwMR--RG--I--lRiDV-NR---N-Ss-NAVnCiLQ-KM-n-AKFMv-TT-NIV- | | |
| TMEBeAn | SPTHYRQTSYTSPTITSVDGWVTWVKLTPLTYPSTPTNSDILTIVSAGDDFTLMP.ISPTKW | | |
| TMEGd7 | -----Q----- | | |
| TMEGd7b | -----Q----- | | |
| TMEDa | -----A--A-----Q-----A-V----- | | |
| Vilyuisk | --S-----S-AA--L--Q----F-ANV-PS-----N----- | | |
| EMCBd | ----F-MVGTDQVN--N-----Q-----P-C-SAK--M--K-S-K--.--AP- | | |
| EMCBc | ----F-MVGTDQ-----Q----- | | |
| EMCDd | ----F-MVGTDQ-----Q----- | | |
| EMCDc | ----F-MVGTDQ-----Q----- | | |
| EMCDv1 | ----F-MVGTDQ-----Q----- | | |
| EMCR | ----F-MVGTDQ-----A-----Q----- | | |
| MengoM | ----F-MVGTDQA-----Q----- | | |
| Mengo37a | ----F-MVGTDL-----A-----Q----- | | |

Serological assay indicating relationship between the Ljungan viruses and diabetes mellitus and myocarditis.

A serological assay using indirect immunofluorescence test using virus infected acetone fixed green monkey kidney cells was established. Patient sera were screened at a 1:8 dilution and positive sera titrated. Sera with a titer of 1:32 or more were considered positive.

Sera from 59 children (age 1-16) from the Stockholm area recently diagnosed with Diabetes Mellitus (DM) and 34 control children from the same geographic area, were tested for presence of antibodies to the three viruses of the invention. Nineteen of the 59 (32%) DM patients screened positive and 2 of the 34 (6%) controls were found positive to one or more of the 3 viruses (significant difference $p=0.002$, Fisher's exact test). Nine recently diagnosed DM patients (age 23-46) from Medelpad county were also tested. Two controls were selected for each adult DM patient and they were matched for age, sex and geographic area of residence. Five of the nine (56%) DM patients and one of the 18 (6%) control patients were found positive to one or more of the 3 viruses (significant difference $p=0.008$ Fisher's exact test).

Serum was also available from 5 athletes dying suddenly in myocarditis. Three controls were selected for each myocarditis patient and they were matched for age, sex and geographic area of residence. Four of the 5 (80%) patients dying from myocarditis and 1 of the 15 (7%) controls were found positive to one or more of the three Ljungan viruses (significant difference $p=0.005$, Fisher's exact test).

DESCRIPTION OF DIFFERENT ASPECTS OF THE INVENTION

In the following, different aspects of the invention will be disclosed. However, all of these aspects are related to a new group of picornaviruses.

Thus, a first aspect of the invention is directed to a new group of picornaviruses, namely picornaviruses comprising in their viral genome, more precisely in the non-coding region, a nucleotide sequence corresponding to a cDNA sequence
5 selected from the group consisting of

SEQ ID NO: 1 (Ljungan 87-012)

10 AGTCTAGTCT TATCTTGTAT GTGTCCTGCA CTGAACTTGT TTCTGTCTCT 50
GGAGTGCTCT ACACTTCAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC 100
ACAGGAATCT GCACAGGTGG CTTTCACCTC TGGACAGTGC ATTCCACACC 150
CGCTCCACGG TAGAAGATGA TGTGTGTCTT TGCTTGTGAA AAGCTTGTGA 200
AAATCGTGTG TAGGCGTAGC GGCTACTTGA GTGCCAGCGG ATTACCCCTA 250
GTGGTAACAC TAGC

15

and homologous sequences having at least 75 % homology to the
SEQ ID NO: 1. The picornaviruses of the invention should further cause
mammalian disease.

20 In a preferred embodiment of this aspect of the invention said homologous
sequences have at least 80%, at least 85% or at least 90% homology to the
SEQ ID NO: 1.

In a particularly preferred embodiment said homologous sequence is one of

25

SEQ ID NO: 2 (Ljungan 174F)

30 AGTCTAGTTT CATTCTGTGT GTGTTTGGCA CTGAAATTAT TTCTGTCTCT 50
GGGGTGCTTT ACACTTCAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC 100
ACAGGAATNT GCACAGGTGG CTTTCACCTC TGGACAGTGC ATTCCACACC 150
CGCTCCACAG TAGAAGATGA TGTGTGTCTT TGCTTGTGAA AAGCTTGTGA 200
AAATCGTGTG TAGGCGTAGC GGNTACTTGA GTGCCAGCGG ACNACCCCTA 250
GTGGTAACAC TAGC

and

SEQ ID NO:3 (Ljungan 145SL).

5 AGTTTGGTTC TCTCTTGAGT GTGTTTTGTG TTAGCATAAT TTCTGTCTCT 50
AGAGTGCTTT AACTCTAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC 100
ACAGGAATCT GCACAGGTGG CTTTCACCTC TGGACAGTGC ATTCCATACC 150
CGCTCCACAA TAGAAGATGA TGTATATCTT TGTTTGTGAA ATGCTCATGA 200
AACGTGTGTG TAGGCGTAGC GGCTACTTGA ATGCCAGCGG AACCCCCCTA 250
GTGGTAACAC TAGC.

10

These sequences (ID NO: 2 and 3) have 94% homology and 91% homology to the SEQ ID NO: 1, respectively.

15

It should be understood that homologies in the coding region of different viruses of the invention may vary considerably, but in the non-coding region they share a homology of at least 75% with the SEQ ID NO: 1.

20

The nucleotide sequences, SEQ ID NO: 1, 2 and 3, correspond to approximately nucleotides 557 - 808 (a conserved region) in the genome of encephalomyelitis virus (EMCV). These three viruses have been isolated from wild rodents, more precisely bank voles. The viruses can be multiplied in cell lines, and for a large-scale production of picornavirus products the virus genome can be inserted into other microorganisms.

25

A second aspect of the invention is directed to a protein comprising an amino acid sequence selected from the group consisting of

SEQ ID NO: 4 (partial structural protein of Ljungan 145)

| | | | | | | | | | | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | Lys | Asp | Leu | Met | Glu | Ile | Ala | Arg | Met | Pro | Ser | Val | Tyr | Lys | Gly | Glu |
| | | | | | 5 | | | | | 10 | | | | | 15 | |
| | Arg | Thr | Glu | Pro | Gly | Gly | Thr | Asn | Gly | Tyr | Phe | Gln | Trp | Ser | His | Thr |
| | | | | 20 | | | | | 25 | | | | | 30 | | |
| 5 | His | Ser | Pro | Ile | Asn | Trp | Val | Phe | Asp | Gly | Gly | Ile | His | Leu | Glu | Asp |
| | | | 35 | | | | | 40 | | | | | 45 | | | |
| | Met | Pro | Asn | Leu | Asn | Leu | Phe | Ser | Ser | Cys | Tyr | Asn | Tyr | Trp | Arg | Gly |
| | | 50 | | | | | 55 | | | | | 60 | | | | |
| | Ser | Thr | Val | Leu | Lys | Leu | Thr | Val | Tyr | Ala | Ser | Thr | Phe | Asn | Lys | Gly |
| 10 | | 65 | | | | 70 | | | | 75 | | | | | | 80 |
| | Arg | Leu | Arg | Met | Ala | Phe | Phe | Pro | Ile | Met | Met | Gln | Gly | Thr | Gln | Arg |
| | | | | | 85 | | | | | 90 | | | | | 95 | |
| | Lys | Lys | His | Lys | Cys | Leu | Phe | Met | Val | Cys | Asp | Ile | Gly | Leu | Asn | Asn |
| | | | | 100 | | | | | 105 | | | | | 110 | | |
| 15 | Thr | Phe | Glu | Met | Thr | Ile | Pro | Tyr | Thr | Trp | Gly | Asn | Trp | Met | Arg | Pro |
| | | | 115 | | | | | 120 | | | | | | 125 | | |
| | Thr | Arg | Gly | Ser | Val | Ile | Gly | Trp | Leu | Arg | Ile | Asp | Val | Leu | Asn | Arg |
| | | 130 | | | | | 135 | | | | | 140 | | | | |
| | Leu | Thr | Tyr | Asn | Ser | Ser | Ser | Pro | Asn | Ala | Val | Asn | Cys | Ile | Leu | Gln |
| 20 | | 145 | | | | 150 | | | | 155 | | | | | | 160 |
| | Val | Lys | Met | Gly | Asn | Asp | Ala | Lys | Phe | Met | Val | Pro | Thr | Thr | Ser | Asn |
| | | | | | 165 | | | | | 170 | | | | | | 175 |

Ile Val Trp ,

and homologous sequences having at least 75% homology to the

25 SEQ ID NO: 4,

and antigenic fragments of the sequences.

In an embodiment of the invention the homologous sequences have at least 80%, at least 85% or at least 90% homology to the SEQ ID NO: 4.

30 The SEQ ID NO: 4 is the result of preliminary partial sequencing of the cDNA sequence from the polyprotein coding sequence of the virus Ljungan 145 SL isolate. Said protein comprising said amino acid sequence SEQ ID NO: 4, said homologous sequences and said antigenic fragments are useful as active ingredients in medicines and as diagnostic reagents in diagnostic kits.

A third aspect of the invention concerns an antiserum or antibody directed against a structural protein of the virus defined in the first aspect of the invention. An example of such a structural protein is defined in the second aspect of the invention. Such an antiserum or antibody is useful as an active ingredient in medicines and as diagnostic reagent in diagnostic kits. Both polyclonal and monoclonal antibodies may be used, and these are suitably produced by using said virus or fragments thereof specific for said virus for immunizing mammals.

A fourth aspect of the invention is directed to an antigen comprising at least a part of a structural protein of the picornavirus defined in the first aspect of the invention, including a subunit thereof. An example of such an antigen is the protein and antigenic parts thereof defined in the second aspect of the invention. Such an antigen of the invention is useful as an active ingredient in medicines and as a diagnostic reagent in diagnostic kits.

A fifth aspect of the invention is directed to a diagnostic kit comprising at least one member from the group consisting of

- a) an antiserum or antibody according to the third aspect of the invention or an antigen-binding part thereof,
- b) an antigen according to the fourth aspect of the invention or an antibody-binding part thereof,
- c) one or several probes designed with respect to the genome of the virus according to the first aspect of the invention,
- and
- d) one or several primers designed with respect to the genome of the virus according to the first aspect of the invention.

The different members of a diagnostic kit will depend on the actual diagnostic method to be used. In addition to the above-listed possible members of the diagnostic kit, said kit may contain positive reference samples, negative

reference samples, diluents, washing solutions and buffers as appropriate. The kit will further be accompanied by instructions for use.

5 The above-listed members a) and b) find use in immunodiagnostic methods, such as enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) or immunofluorescence assay (IFA).

The above-listed members c) and d) find use in direct virus detection. Preferably, a diagnostic method based on the PCR (polymer chain reaction) 10 technique with such primers is utilized in the direct detection of a virus according to the invention.

All of the above mentioned diagnostic methods are well known in the art, and a man of ordinary skill in the art will readily select useful members for a diagnostic 15 kit in relation to the diagnostic method to be used.

A sixth aspect of the invention relates to a vaccine having as an immunizing or neutralizing component a member selected from the group consisting of
20 a) the virus according to the first aspect of the invention,
b) the virus according to the first aspect of the invention in attenuated form,
c) the virus according to the first aspect of the invention in killed form,
d) an antigen according to the fourth aspect of the invention, including a subunit of the virus according to the first aspect of the invention,
and
25 e) DNA corresponding to the genomic RNA of the virus according to the first aspect of the invention.

In an embodiment of this aspect of the invention said vaccine may additionally comprises an adjuvant. Such an adjuvant must of course be an adjuvant which is approved for use in vaccines by authorities responsible for veterinary or 30 human medicines.

The vaccine may contain other ingredients which are needed for specific preparations intended for oral, subcutaneous, intramuscular or intradermal administration. Suitable additional ingredients are disclosed in the European or US Pharmacopoeia.

5

The alternative members a), b) and c) are all examples of conventional whole virus, attenuated virus, and subunit vaccines developed for other types of viruses, and the member d) represents DNA incorporation into body-specific cells, which then will express virus-specific structures and elicit immunity against said virus.

10

A seventh aspect of the invention is directed to a picornavirus according to the first aspect of the invention, optionally in attenuated or killed form, an antiserum or antibody according to the third aspect of the invention or an antigen according to the fourth aspect of the invention, for use in a medicament (for veterinary or human use). An example of such a medicament is a vaccine according to the invention disclosed in the sixth aspect thereof.

15

The eighth aspect of the invention concerns use of a picornavirus according to the first aspect of the invention, optionally in attenuated or killed form, an antiserum or antibody according to the third aspect of the invention or an antigen according to the fourth aspect of the invention, in the preparation of a medicament for prophylactic or therapeutic treatment of a disease caused by said virus.

20

25

In an embodiment of said use the disease caused by said virus is one of Myocarditis, Cardiomyopathia, Guillain Barré Syndrome, and Diabetes Mellitus, Multiple Sclerosis, Chronic Fatigue Syndrome, Myasthenia Gravis, Amyotrophic Lateral Sclerosis, Dermatomyositis, Polymyositis, Spontaneous Abortion, and Sudden Infant Death Syndrome.

30

A ninth aspect of the invention is directed to a method of prophylactic or therapeutic treatment of a disease caused by a virus according to the first aspect of the invention in a mammal, including human, which comprises administering to said mammal a prophylactically or therapeutically effective amount of a medicament comprising as an active ingredient a member of the group consisting of

- a) the virus according to the first aspect of the invention,
- b) the virus according to the first aspect of the invention in attenuated form,
- c) the virus according to the first aspect of the invention in killed form,
- d) an antigen according to the fourth aspect of the invention, including a subunit of the virus according to the first aspect of the invention,
- and
- e) DNA corresponding to the genomic RNA of the virus according to the first aspect of the invention.

In an embodiment of said method the disease caused by said virus is one of Myocarditis, Cardiomyopathia, Guillain Barré Syndrome, and Diabetes Mellitus, Multiple Sclerosis, Chronic Fatigue Syndrome, Myasthenia Gravis, Amyotrophic Lateral Sclerosis, Dermatomyositis, Polymyositis, Spontaneous Abortion, and Sudden Infant Death Syndrome.

The actual dosage regimen will be determined by the vaccine producer based on animal experiments and clinical trials.

REFERENCES

1. Wesslén, L. *et al.* Myocarditis caused by *Chlamydia pneumoniae* (TWAR) and sudden unexpected death in a Swedish elite orienteer. The Lancet, **340**, 427-428 (1992).
5
2. Gard, S. Heller L. Hemagglutination by Col-MM-virus. Proc. Soc. Exp. Biol. Med. 76, 68-73 1951.
3. Francki, R.I.B., C.M. Fauquet, D.L. Knudson, and F. Brown (ed). 1991. Classification and nomenclature of viruses. Fifth report of the International Committee in Taxonomy of viruses. Arch. Virol. 1991 (Suppl. 2): 320-326.
10
4. Palmenberg, A.C. 1989. Sequence alignment of picornaviral capsid proteins, p. 211-241. In B.L. Semler and E. Ehrenfeld (ed.), Molecular aspects of picornavirus infection and detection. American Society for Microbiology, Washington. D.C.
15
5. Stanway, G. 1990. Structure, function and evolution of picornaviruses. J. Gen. Virol. 71:2483-2501.
20
6. Lipton, H.L., A. Friedmann, P. Sethi, and J.R. Crowther. 1983. Characterization of Vilyuisk virus as a picornavirus. J. Med. Virol. 12:195-203.
7. Zimmerman, J.J. Encephalomyocarditis. In: Handbook Series of Zoonoses. G. B. Beran (Ed.), CRC, Press Inc., Boca Raton, FL USA (1994).
25
8. Hubbard, G.B. *et al.* An encephalomyocarditis virus epizootic in a baboon colony. *Lab. Anim. Sci.* **42**, 223-239 (1992).
30
9. Gaskin, J.M. *et al.* The tragedy of encephalomyocarditis virus infection in zoological parks in Florida. *Proc. Am. Assoc. Zoo. Vet* 1-7 (1980).

10. Craighead, J.E & McLane, M.F. Diabetes Mellitus: induction in mice by encephalomyocarditis virus. *Science* **162**, 913-915 (1968).
- 5 11. Hayashi, K., Boucher, D.W. & Notkins, A.L. Virus induced diabetes mellitus. II. Relationship between β -cell damage and hyperglycemia in mice infected with encephalomyocarditis virus. *Am. J. Pathol.* **75**, 91-102 (1974).
- 10 12. Dal Canto, M.C. Experimental models of virus-induced demyelination, in Handbook of Multiple Sclerosis, in Cook SD (ed), New York, Marcel Decker, pp 63-100 (1990).
- 15 13. Hirasawa, K., Han, J. Takeda, M., Itagaki, S. & Doi, K. J. Encephalomyocarditis (EMC) virus induced myocarditis by different virus variants and mouse strains. *Vet. Med. Sci.* **54**, 1125-1129 (1992).
14. Levin, R. *et al.* EMC virus infection in baboons as a model for studies on antiviral substances. *Antiviral Research* **6**, 277-283 (1986).
- 20 15. Blanchard, J.L., Soike, K. & Baskin, G.B. Encephalomyocarditis virus infection in African green and squirrel monkeys: Comparison of pathologic effect. *Laboratory Animal Science* **37**, 635-639 (1987).
- 25 16. Lipton, H.L. & Dal Canto, M.C. Theiler's virus induced demyelination: prevention by immunosuppression. *Science* **192**, 62-64 (1976).
17. Gajdusek, D.C. Encephalomyocarditis virus infection in childhood. *Pediatrics* **16**, 902-906 (1955).
- 30 18. Niklasson, B. & Le Duc, J. Epidemiology of nephropathia epidemica in Sweden. *J. Inf. Dis.* **155**:269-276 (1987).

19. Riggs, J.L.: Immunofluorescent staining. In: Diagnostic procedures for Viral, Rickettsial, and Chlamydial infections, Am. Public. Health Assoc., Washington 1979, 5th ed., p. 141.
- 5 20. Earley, E., Peralta, P.H. & Johnson, K.M.: A plaque neutralization method for arboviruses. Proc. Soc. Exp. Biol. Med. 125:741. (1967)

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: BO NIKLASSON
- (B) STREET: Sibyllegatan 15
- (C) CITY: Stockholm
- (E) COUNTRY: Sweden
- (F) POSTAL CODE (ZIP): 114 42

(ii) TITLE OF INVENTION: NEW PICORNAVIRUSES, VACCINES AND
DIAGNOSTIC KITS.

(iii) NUMBER OF SEQUENCES: 4

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: PICORNAVIRIDAE
- (C) INDIVIDUAL ISOLATE: LJUNGAN 87-012

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```
AGTCTAGTCT TATCTTGTAT GTGTCCTGCA CTGAACTTGT TTCTGTCTCT GGAGTGCTCT    60
ACACTTCAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC ACAGGAATCT GCACAGGTGG    120
CTTTCACCTC TGGACAGTGC ATTCCACACC CGCTCCACGG TAGAAGATGA TGTGTGTCTT    180
TGCTTGTGAA AAGCTTGTGA AAATCGTGTG TAGGCGTAGC GGCTACTTGA GTGCCAGCGG    240
ATTACCCCTA GTGGTAACAC TAGC                                           264
```

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 261 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Picornaviridae
- (C) INDIVIDUAL ISOLATE: Ljungan 174F

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

```
AGTCTAGTTT CATTCTGTGT GTGTTTGGCA CTGAAATTAT TTCTGTCTCT GGGGTGCTTT    60
ACACTTCAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC ACAGGAATTG CACAGGTGGC    120
TTTCACCTCT GGACAGTGCA TTCCACACCC GCTCCACAGT AGAAGATGAT GTGTGTCTTT    180
GCTTGTGAAA AGCTTGTGAA AATCGTGTGT AGGCGTAGCG GTACTTGAGT GCCAGCGGAC    240
ACCCCTAGTG GTAACACTAG C                                                261
```

2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Picornaviridae
- (C) INDIVIDUAL ISOLATE: Ljungan 145SL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

```
AGTTTGTTTC TCTCTTGAGT GTGTTTTGTG TTAGCATAAT TTCTGTCTCT AGAGTGCTTT    60
ACACTCTAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC ACAGGAATCT GCACAGGTG    120
CTTTCACCTC TGGACAGTGC ATTCCATACC CGCTCCACAA TAGAAGATGA TGTATATCTT    180
TGTTTGTGAA ATGCTCATGA AACGTGTGTG TAGGCGTAGC GGCTACTTGA ATGCCAGCGG    240
AACCCCCCTA GTGGTAACAC TAGC                                                264
```

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 179 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Picornaviridae
- (C) INDIVIDUAL ISOLATE: Ljungan 145SL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

```

Lys Asp Leu Met Glu Ile Ala Arg Met Pro Ser Val Tyr Lys Gly Glu
1           5           10           15

Arg Thr Glu Pro Gly Gly Thr Asn Gly Tyr Phe Gln Trp Ser His Thr
          20           25           30

His Ser Pro Ile Asn Trp Val Phe Asp Gly Gly Ile His Leu Glu Asp
          35           40           45

Met Pro Asn Leu Asn Leu Phe Ser Ser Cys Tyr Asn Tyr Trp Arg Gly
          50           55           60

Ser Thr Val Leu Lys Leu Thr Val Tyr Ala Ser Thr Phe Asn Lys Gly
          65           70           75           80

Arg Leu Arg Met Ala Phe Phe Pro Ile Met Met Gln Gly Thr Gln Arg
          85           90           95

Lys Lys His Lys Cys Leu Phe Met Val Cys Asp Ile Gly Leu Asn Asn
          100          105          110

Thr Phe Glu Met Thr Ile Pro Tyr Thr Trp Gly Asn Trp Met Arg Pro
          115          120          125

Thr Arg Gly Ser Val Ile Gly Trp Leu Arg Ile Asp Val Leu Asn Arg
          130          135          140

Leu Thr Tyr Asn Ser Ser Ser Pro Asn Ala Val Asn Cys Ile Leu Gln
          145          150          155          160

Val Lys Met Gly Asn Asp Ala Lys Phe Met Val Pro Thr Thr Ser Asn
          165          170          175

Ile Val Trp

```

22 -10- 1998

28

29140/BN

CLAIMS

5

1. Ljungan picornavirus, comprising in the non-coding region of its viral genome a nucleotide sequence corresponding to a cDNA sequence selected from the group consisting of

10

SEQ ID NO: 1 (Ljungan 87-012)

15

```

AGTCTAGTCT TATCTTGTAT GTGTCCTGCA CTGAACTTGT TTCTGTCTCT 50
GGAGTGCTCT ACACTTCAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC 100
ACAGGAATCT GCACAGGTGG CTTTCACCTC TGGACAGTGC ATTCCACACC 150
CGCTCCACGG TAGAAGATGA TGTGTGTCTT TGCTTGTGAA AAGCTTGTGA 200
AAATCGTGTG TAGGCGTAGC GGCTACTTGA GTGCCAGCGG ATTACCCCTA 250
GTGGTAACAC TAGC

```

20

and homologous sequences having at least 75% homology to the SEQ ID NO: 1,
and further, causing mammalian disease.

25

2. Ljungan picornavirus according to claim 1, wherein said homologous sequences have at least 80%, at least 85% or at least 90% homology to the SEQ ID NO: 1.

30

3. Ljungan picornavirus according to claim 2, wherein said homologous sequence is one of

35

```

AGTCTAGTTT CATTCTGTGT GTGTTTGGCA CTGAAATTAT TTCTGTCTCT 50
GGGGTGCTTT ACACTTCAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC 100
ACAGGAATNT GCACAGGTGG CTTTCACCTC TGGACAGTGC ATTCCACACC 150
CGCTCCACAG TAGAAGATGA TGTGTGTCTT TGCTTGTGAA AAGCTTGTGA 200
AAATCGTGTG TAGGCGTAGC GGNTACTTGA GTGCCAGCGG ACNACCCCTA 250
GTGGTAACAC TAGC

```

40

and
SEQ ID NO:3 (Ljungan 145SL).

22 -10- 1998

29

AGTTTGGTTC TCTCTTGAGT GTGTTTGTG TTAGCATAAT TTCTGTCTCT 50
AGAGTGCTTT ACACTCTAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC 100
ACAGGAATCT GCACAGGTGG CTTTCACCTC TGGACAGTGC ATTCCATACC 150
CGCTCCACAA TAGAAGATGA TGTATATCTT TGTTTGTGAA ATGCTCATGA 200
5 AACGTGTGTG TAGGCGTAGC GGCTACTTGA ATGCCAGCGG AACCCCCCTA 250
GTGGTAACAC TAGC.

4. Protein comprising an amino acid sequence selected from the group
consisting of
SEQ ID NO: 4 (partial structural protein of Ljungan 145SL)

| | | | | | | | | | | | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|
| | Lys | Asp | Leu | Met | Glu | Ile | Ala | Arg | Met | Pro | Ser | Val | Tyr | Lys | Gly | Glu | |
| | | | | | 5 | | | | | 10 | | | | | 15 | | |
| 15 | Arg | Thr | Glu | Pro | Gly | Gly | Thr | Asn | Gly | Tyr | Phe | Gln | Trp | Ser | His | Thr | |
| | | | | 20 | | | | | 25 | | | | | 30 | | | |
| | His | Ser | Pro | Ile | Asn | Trp | Val | Phe | Asp | Gly | Gly | Ile | His | Leu | Glu | Asp | |
| | | | | 35 | | | | 40 | | | | | 45 | | | | |
| | Met | Pro | Asn | Leu | Asn | Leu | Phe | Ser | Ser | Cys | Tyr | Asn | Tyr | Trp | Arg | Gly | |
| 20 | | 50 | | | | | 55 | | | | | 60 | | | | | |
| | Ser | Thr | Val | Leu | Lys | Leu | Thr | Val | Tyr | Ala | Ser | Thr | Phe | Asn | Lys | Gly | |
| | 65 | | | | | 70 | | | | | 75 | | | | 80 | | |
| | Arg | Leu | Arg | Met | Ala | Phe | Phe | Pro | Ile | Met | Met | Gln | Gly | Thr | Gln | Arg | |
| | | | | 85 | | | | | 90 | | | | | 95 | | | |
| 25 | Lys | Lys | His | Lys | Cys | Leu | Phe | Met | Val | Cys | Asp | Ile | Gly | Leu | Asn | Asn | |
| | | | | 100 | | | | | 105 | | | | | 110 | | | |
| | Thr | Phe | Glu | Met | Thr | Ile | Pro | Tyr | Thr | Trp | Gly | Asn | Trp | Met | Arg | Pro | |
| | | | | 115 | | | | | 120 | | | | | 125 | | | |
| | Thr | Arg | Gly | Ser | Val | Ile | Gly | Trp | Leu | Arg | Ile | Asp | Val | Leu | Asn | Arg | |
| 30 | | 130 | | | | | 135 | | | | | 140 | | | | | |
| | Leu | Thr | Tyr | Asn | Ser | Ser | Ser | Pro | Asn | Ala | Val | Asn | Cys | Ile | Leu | Gln | |
| | 145 | | | | 150 | | | | | 155 | | | | 160 | | | |
| | Val | Lys | Met | Gly | Asn | Asp | Ala | Lys | Phe | Met | Val | Pro | Thr | Thr | Ser | Asn | |
| | | | | 165 | | | | | 170 | | | | | 175 | | | |
| 35 | Ile | Val | Trp | | | | | | | | | | | | | | |

and homologous sequences having at least 75% homology to the SEQ ID
NO: 4,
and antigenic fragments of the sequences.

22-10-1998

30

5. Antiserum or antibody directed against a structural protein of the virus according to any one of claims 1-3.

5 6. Antigen comprising at least a part of a structural protein of the picornavirus according to any one of claims 1-3.

7. Diagnostic kit comprising at least one member from the group consisting of an antiserum or antibody according to claim 5 or an antigen-binding part thereof,
10 an antigen according to claim 6 or an antibody-binding part thereof,
one or several probes designed with respect to the genome of the virus according to any one of claims 1-3,
and
one or several primers designed with respect to the genome of the virus
15 according to any one of claims 1-3.

8. Vaccine having as an immunizing or neutralizing component a member selected from the group consisting of
a) the virus according to any one of claims 1-3,
20 b) the virus according to any one of claims 1-3 in attenuated form,
c) the virus according to any one of claims 1-3 in killed form,
d) an antigen according to claim 6, including a subunit of the virus according to any one of claims 1-3,
and
25 e) DNA corresponding to the genomic RNA of the virus according to any one of claims 1-3.

9. Vaccine according to claim 8 which additionally comprises an adjuvant.

30 10. Ljungan picornavirus according to any one of the claims 1-3, optionally in attenuated or killed form, an antiserum or antibody according to claim 5 or an antigen according to claim 6 for use in a medicament.

11. Use of a Ljungan picornavirus according to any one of the claims 1-3, optionally in attenuated or killed form, an antiserum or antibody according to claim 5 or an antigen according to claim 6, in the preparation of a medicament for prophylactic or therapeutic treatment of a disease caused by said virus.

5

12. Use according to claim 11, wherein the disease caused by said virus is one of Myocarditis, Cardiomyopathia, Guillain Barré Syndrome, and Diabetes Mellitus, Multiple Sclerosis, Chronic Fatigue Syndrome, Myasthenia Gravis, Amyotrophic Lateral Sclerosis, Dermatomyositis, Polymyositis, Spontaneous Abortion, and Sudden Infant Death Syndrome.

10

13. Method of prophylactic or therapeutic treatment of a disease caused by a virus according to any one of the claims 1-3 in a mammal, including human, which comprises administering to said mammal a prophylactically or therapeutically effective amount of a medicament comprising as an active ingredient a member of the group consisting of

15

a) the virus according to any one of claims 1-3,

b) the virus according to any one of claims 1-3 in attenuated form,

c) the virus according to any one of claims 1-3 in killed form,

20

d) an antigen according to claim 6, including a subunit of the virus according to any one of claims 1-3,

and

e) DNA corresponding to the genomic RNA of the virus according to any one of claims 1-3.

25

14. Method according to claim 13, wherein the disease caused by said virus is one of Myocarditis, Cardiomyopathia, Guillain Barré Syndrome, and Diabetes Mellitus, Multiple Sclerosis, Chronic Fatigue Syndrome, Myasthenia Gravis, Amyotrophic Lateral Sclerosis, Dermatomyositis, Polymyositis, Spontaneous Abortion, and Sudden Infant Death Syndrome.

30

CLAIMS

5

1. Picornavirus, comprising in the non-coding region of its viral genome a nucleotide sequence corresponding to a cDNA sequence selected from the group consisting of

10

SEQ ID NO: 1 (Ljungan 87-012)

15

```
AGTCTAGTCT TATCTTGTAT GTGTCCTGCA CTGAACTTGT TTCTGTCTCT    50
GGAGTGCTCT ACACTTCAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC    100
ACAGGAATCT GCACAGGTGG CTTTCACCTC TGGACAGTGC ATTCCACACC    150
CGCTCCACGG TAGAAGATGA TGTGTGTCTT TGCTTGTGAA AAGCTTGTGA    200
AAATCGTGTG TAGGCGTAGC GGCTACTTGA GTGCCAGCGG ATTACCCCTA    250
GTGGTAACAC TAGC
```

20

and homologous sequences having at least 75% homology to the SEQ ID NO: 1,
and further, causing mammalian disease.

2. Picornavirus according to claim 1, wherein said homologous sequences have at least 80%, at least 85% or at least 90% homology to the SEQ ID NO: 1.

25

3. Picornavirus according to claim 2, wherein said homologous sequence is one of

30

SEQ ID NO:2 (Ljungan 174F)

35

```
AGTCTAGTTT CATTCTGTGT GTGTTTGGCA CTGAAATTAT TTCTGTCTCT    50
GGGGTGCTTT ACACTTCAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC    100
ACAGGAATNT GCACAGGTGG CTTTCACCTC TGGACAGTGC ATTCCACACC    150
CGCTCCACAG TAGAAGATGA TGTGTGTCTT TGCTTGTGAA AAGCTTGTGA    200
AAATCGTGTG TAGGCGTAGC GGNTACTTGA GTGCCAGCGG ACNACCCCTA    250
GTGGTAACAC TAGC
```

and

SEQ ID NO:3 (Ljungan 145SL).

40

AGTTTGGTTC TCTCTTGAGT GTGTTTTGTG TTAGCATAAT TTCTGTCTCT 50
 AGAGTGCTTT ACACTCTAGT AGGGGCTGTA CCCGGGCGGT CCCACTCTTC 100
 ACAGGAATCT GCACAGGTGG CTTTCACCTC TGGACAGTGC ATTCCATACC 150
 CGCTCCACAA TAGAAGATGA TGTATATCTT TGTGTGTGAA ATGCTCATGA 200
 AACGTGTGTG TAGGCGTAGC GGCTACTTGA ATGCCAGCGG AACCCCCCTA 250
 GTGGTAACAC TAGC.

4. Protein comprising an amino acid sequence selected from the group
 consisting of
 SEQ ID NO: 4 (partial structural protein of Ljungan 145SL)

| | | | | | | | | | | | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|
| | Lys | Asp | Leu | Met | Glu | Ile | Ala | Arg | Met | Pro | Ser | Val | Tyr | Lys | Gly | Glu | |
| | | | | | 5 | | | | | 10 | | | | | 15 | | |
| 15 | Arg | Thr | Glu | Pro | Gly | Gly | Thr | Asn | Gly | Tyr | Phe | Gln | Trp | Ser | His | Thr | |
| | | | | 20 | | | | | 25 | | | | | 30 | | | |
| | His | Ser | Pro | Ile | Asn | Trp | Val | Phe | Asp | Gly | Gly | Ile | His | Leu | Glu | Asp | |
| | | | | 35 | | | | 40 | | | | | 45 | | | | |
| | Met | Pro | Asn | Leu | Asn | Leu | Phe | Ser | Ser | Cys | Tyr | Asn | Tyr | Trp | Arg | Gly | |
| 20 | | 50 | | | | | 55 | | | | | 60 | | | | | |
| | Ser | Thr | Val | Leu | Lys | Leu | Thr | Val | Tyr | Ala | Ser | Thr | Phe | Asn | Lys | Gly | |
| | 65 | | | | | 70 | | | | | 75 | | | | | 80 | |
| | Arg | Leu | Arg | Met | Ala | Phe | Phe | Pro | Ile | Met | Met | Gln | Gly | Thr | Gln | Arg | |
| | | | | 85 | | | | | 90 | | | | | 95 | | | |
| 25 | Lys | Lys | His | Lys | Cys | Leu | Phe | Met | Val | Cys | Asp | Ile | Gly | Leu | Asn | Asn | |
| | | | | 100 | | | | | 105 | | | | | 110 | | | |
| | Thr | Phe | Glu | Met | Thr | Ile | Pro | Tyr | Thr | Trp | Gly | Asn | Trp | Met | Arg | Pro | |
| | | | | 115 | | | | | 120 | | | | 125 | | | | |
| | Thr | Arg | Gly | Ser | Val | Ile | Gly | Trp | Leu | Arg | Ile | Asp | Val | Leu | Asn | Arg | |
| 30 | | 130 | | | | | 135 | | | | | 140 | | | | | |
| | Leu | Thr | Tyr | Asn | Ser | Ser | Ser | Pro | Asn | Ala | Val | Asn | Cys | Ile | Leu | Gln | |
| | 145 | | | | | 150 | | | | | 155 | | | | | 160 | |
| | Val | Lys | Met | Gly | Asn | Asp | Ala | Lys | Phe | Met | Val | Pro | Thr | Thr | Ser | Asn | |
| | | | | 165 | | | | | 170 | | | | | 175 | | | |
| 35 | Ile | Val | Trp | , | | | | | | | | | | | | | |

and homologous sequences having at least 75 % homology to the SEQ ID
 NO: 4,
 and antigenic fragments of the sequences.

5. Antiserum or antibody directed against a structural protein of the virus according to any one of claims 1-3.
- 5 6. Antigen comprising at least a part of a structural protein of the picornavirus according to any one of claims 1-3.
7. Diagnostic kit comprising at least one member from the group consisting of an antiserum or antibody according to claim 5 or an antigen-binding part thereof,
10 an antigen according to claim 6 or an antibody-binding part thereof,
one or several probes designed with respect to the genome of the virus according to any one of claims 1-3,
and
one or several primers designed with respect to the genome of the virus
15 according to any one of claims 1-3.
8. Vaccine having as an immunizing or neutralizing component a member selected from the group consisting of
a) the virus according to any one of claims 1-3,
20 b) the virus according to any one of claims 1-3 in attenuated form,
c) the virus according to any one of claims 1-3 in killed form,
d) an antigen according to claim 6, including a subunit of the virus according to any one of claims 1-3,
and
25 e) DNA corresponding to the genomic RNA of the virus according to any one of claims 1-3.
9. Vaccine according to claim 8 which additionally comprises an adjuvant.
- 30 10. Picornavirus according to any one of the claims 1-3, optionally in attenuated or killed form, an antiserum or antibody according to claim 5 or an antigen according to claim 6 for use in a medicament.

11. Use of a picornavirus according to any one of the claims 1-3, optionally in attenuated or killed form, an antiserum or antibody according to claim 5 or an antigen according to claim 6, in the preparation of a medicament for prophylactic or therapeutic treatment of a disease caused by said virus.

12. Use according to claim 11, wherein the disease caused by said virus is one of Myocarditis, Cardiomyopathia, Guillain Barré Syndrome, and Diabetes Mellitus, Multiple Sclerosis, Chronic Fatigue Syndrome, Myasthenia Gravis, Amyotrophic Lateral Sclerosis, Dermatomyositis, Polymyositis, Spontaneous Abortion, and Sudden Infant Death Syndrome.

13. Method of prophylactic or therapeutic treatment of a disease caused by a virus according to any one of the claims 1-3 in a mammal, including human, which comprises administering to said mammal a prophylactically or therapeutically effective amount of a medicament comprising as an active ingredient a member of the group consisting of

- a) the virus according to any one of claims 1-3,
- b) the virus according to any one of claims 1-3 in attenuated form,
- c) the virus according to any one of claims 1-3 in killed form,
- d) an antigen according to claim 6, including a subunit of the virus according to any one of claims 1-3,

and

- e) DNA corresponding to the genomic RNA of the virus according to any one of claims 1-3.

14. Method according to claim 13, wherein the disease caused by said virus is one of Myocarditis, Cardiomyopathia, Guillain Barré Syndrome, and Diabetes Mellitus, Multiple Sclerosis, Chronic Fatigue Syndrome, Myasthenia Gravis, Amyotrophic Lateral Sclerosis, Dermatomyositis, Polymyositis, Spontaneous Abortion, and Sudden Infant Death Syndrome.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/SE 97/01515

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| US 5229111 A | 20/07/93 | NONE | |

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/01515

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07K 14/085, A61K 39/125, G01N 33/569
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07K, A61K, G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim N . |
|-----------|--|-----------------------|
| A | Proc. Natl. Acad. Sci., Volume 89, Sept 1992, T Hyypiä et al, "A distinct picornavirus group identified by sequence analysis", page 8847 - page 8851, see figure 2, positions 1770-2330 | 1-4 |
| Y | -- | 5-12 |
| Y | Journal of General Virology, Volume 76, 1995, Hee Sook Jun et al, "Cloning and expression of the VPI major capsid protein of diabetogenic encephalomyocarditis (EMC) virus and prevention of EMC virus-induced diabetes by immunization with the recombinant VPI protein" page 2557 - page 2566 | 5-12 |
| | -- | |

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

- * Special categories of cited documents:
- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed
- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

| | |
|--|---|
| Date of the actual completion of the international search | Date of mailing of the international search report |
| 16 December 1997 | 02 -01- 1998 |
| Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile N . +46 8 666 02 86 | Authorized officer Carl-Olof Gustafsson Telephone No. +46 8 782 25 00 |

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---|-----------------------|
| Y | Exp. Anim., Volume 44, No 3, 1995, K Dan et al, "Characterization of Insulin-Dependent Diabetes Mellitus Induced by a New Variant (DK-27) of Encephalomyocarditis Virus in DBA/2 Mice" page 211 - page 218 -- | 5-12 |
| A | Proc. Soc. Exp. Biol. Med., Volume 205, No 2, 1994, M Tolbert et al, "Persistence of the D Variant of Encephalomyocarditis Virus in the ICR-Swiss Mouse (43687)" page 124 -- | 1-12 |
| A | Dialog Information Services, file 34, SciSearch, Dialog accession no. 14364904, Samuelson A et al: "Sequence-analysis of echoviruses in a major antigenic region eliciting enteroviral cross- -reactive antibodies"; & ARCHIVES OF VIROLOGY, 1995, V140, N11, P2085-2091 -- | 4 |
| Y | Dialog Information Services, file 154, MEDLINE, Dialog accession no. 08478765, Medline accession no. 96050833, Samuelson A et al: "Molecular basis for serological cross-reactivity between entero- viruses"; Clin Diagn Lab Immunol May 1994, 1 (3) p336-41 -- | 5-12 |
| Y | Dialog Information Services, file 154, MEDLINE, Dialog accession no. 08518839, Medline accession no. 96125356, Schultheiss T et al: "Polyprotein processing in echovirus 22: a first assessment"; & Biochem Biophys Res Commun Dec 26 1995, 217 p1120-7 -- | 5-12 |
| A | Dialog Information Services, file 154, MEDLINE, Dialog accession no. 06649814, Medline accession no. 90244381, Collier BA et al: "Echovirus 22 is an atypical enterovirus"; & J Virol Jun 1990, 64 (6) p2692-701 -- | 1-12 |

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/01515

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| A | US 5229111 A (ANN C. PALMENBERG ET AL), 20 July 1993 (20.07.93) ----- | 6,8-12 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/SE 97/01515

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| US 5229111 A | 20/07/93 | NONE | |